



---

# How Much Meson Physics Can One Tie to DCSB?

Peter Tandy  
Center for Nuclear Research  
Kent State University



# Topics

---

- DSE modeling of meson physics (ladder-rainbow)
- Summary: successes and problems
- More recently: heavy quark mesons, DIS on pion
- Steps beyond ladder-rainbow

# Lattice-QCD and DSE-based modeling

---

- Lattice:  $\langle \mathcal{O} \rangle = \int D\bar{q}qG \mathcal{O}(\bar{q}, q, G) e^{-\mathcal{S}[\bar{q}, q, G]}$ 
  - Euclidean metric, x-space, Monte-Carlo
  - Issues: lattice spacing and vol, sea and valence  $m_q$ , fermion Det
  - **Large time limit**  $\Rightarrow$  nearest hadronic mass pole
- EOMs (DSEs):  $0 = \int D\bar{q}qG \frac{\delta}{\delta q(x)} e^{-\mathcal{S}[\bar{q}, q, G] + (\bar{\eta}, q) + (\bar{q}, \eta) + (J, G)}$ 
  - Euclidean metric, p-space, continuum integral eqns
  - Issues: truncation and phenomenology—not full QCD
  - **Analytic contin.**  $\Rightarrow$  nearest hadronic mass pole
  - Can be quick to identify systematics, mechanisms,  $\dots$

# DSE-based modeling of Hadron Physics

---

- Soft physics: truncate DSEs to min: 2-pt, 3-pt fns
- Should be **relativistically covariant**—convenient for decays, Form Factors, etc
  - **No boosts needed on wavefns of recoiling bound st.**
  - **$\infty$  d.o.f.  $\rightarrow$  few quasi-particle effective d.o.f.**
- Do not make a 3-dimensional reduction
- Preserve 1-loop QCD renorm group behavior in UV
- Preserve global symmetries, conserved em currents, etc
- Preserve PCAC  $\Rightarrow$  Goldstone's Thm
- Can't preserve local color gauge covariance—just choose Landau gauge [RG fixed pt]
- Parameterize the deep infrared (large distance) QCD coupling

# Constraints on Modeling

---


- Preserve vector WTI, and **axial vector WTI**

E.g.

$$-iP_\mu \Gamma_{5\mu}(k; P) = S^{-1}(k_+) \gamma_5 \frac{\tau}{2} + \gamma_5 \frac{\tau}{2} S^{-1}(k_-) - 2m_q(\mu) \Gamma_5(k; P)$$

- $\Rightarrow$  kernels of  $DSE_q$  and  $K_{BSE}$  are related
- Ladder-rainbow is the simplest implementation
- **Goldstone Theorem preserved**, ps octet masses good, indep of model details
- **DCSB**  $\Rightarrow \pi$ :  $\Gamma_\pi^0(p^2) = \frac{i\gamma_5}{f_\pi^0} \left[ \frac{1}{4} \text{tr} S_0^{-1}(p^2) \right] + \dots$
- Here, 1-body and 2-body systems are the same


# Ladder-Rainbow Model



- $K_{\text{BSE}} \rightarrow -\gamma_\mu \frac{\lambda^a}{2} 4\pi\alpha_{\text{eff}}(q^2) D_{\mu\nu}^{\text{free}}(q) \gamma_\nu \frac{\lambda^a}{2}$

- $\alpha_{\text{eff}}(q^2) \xrightarrow{IR} \langle \bar{q}q \rangle_{\mu=1} \text{ GeV} = -(240\text{MeV})^3$ , incl vertex dressing

- $\alpha_{\text{eff}}(q^2) \xrightarrow{UV} \alpha_s^{1-\text{loop}}(q^2)$



- P. Maris & P.C. Tandy, PRC60, 055214 (1999)

$M_\rho, M_\phi, M_{K^*}$  good to 5%,  $f_\rho, f_\phi, f_{K^*}$  good to 10%

## Summary of light meson results

$m_{u=d} = 5.5 \text{ MeV}$ ,  $m_s = 125 \text{ MeV}$  at  $\mu = 1 \text{ GeV}$

### Pseudoscalar (PM, Roberts, PRC56, 3369)

	expt.	calc.
$-\langle \bar{q}q \rangle_\mu^0$	$(0.236 \text{ GeV})^3$	$(0.241^\dagger)^3$
$m_\pi$	0.1385 GeV	$0.138^\dagger$
$f_\pi$	0.0924 GeV	$0.093^\dagger$
$m_K$	0.496 GeV	$0.497^\dagger$
$f_K$	0.113 GeV	0.109

### Charge radii (PM, Tandy, PRC62, 055204)

$r_\pi^2$	0.44 fm <sup>2</sup>	0.45
$r_{K^+}^2$	0.34 fm <sup>2</sup>	0.38
$r_{K^0}^2$	-0.054 fm <sup>2</sup>	-0.086

### $\gamma\pi\gamma$ transition (PM, Tandy, PRC65, 045211)

$g_{\pi\gamma\gamma}$	0.50	0.50
$r_{\pi\gamma\gamma}^2$	0.42 fm <sup>2</sup>	0.41

### Weak $K_{l3}$ decay (PM, Ji, PRD64, 014032)

$\lambda_+(e3)$	0.028	0.027
$\Gamma(K_{e3})$	$7.6 \cdot 10^6 \text{ s}^{-1}$	7.38
$\Gamma(K_{\mu3})$	$5.2 \cdot 10^6 \text{ s}^{-1}$	4.90

### Vector mesons

(PM, Tandy, PRC60, 055214)

$m_{\rho/\omega}$	0.770 GeV	0.742
$f_{\rho/\omega}$	0.216 GeV	0.207
$m_{K^*}$	0.892 GeV	0.936
$f_{K^*}$	0.225 GeV	0.241
$m_\phi$	1.020 GeV	1.072
$f_\phi$	0.236 GeV	0.259

### Strong decay (Jarecke, PM, Tandy, PRC67, 035202)

$g_{\rho\pi\pi}$	6.02	5.4
$g_{\phi KK}$	4.64	4.3
$g_{K^*K\pi}$	4.60	4.1

### Radiative decay

(PM, nucl-th/0112022)

$g_{\rho\pi\gamma}/m_\rho$	0.74	0.69
$g_{\omega\pi\gamma}/m_\omega$	2.31	2.07
$(g_{K^*K\gamma}/m_{K^*})^+$	0.83	0.99
$(g_{K^*K\gamma}/m_{K^*})^0$	1.28	1.19

### Scattering length

(PM, Cotanch, PRD66, 116010)

$a_0^0$	0.220	0.170
$a_0^2$	0.044	0.045
$a_1^1$	0.038	0.036

bsampl


# DSE kernel constrained from Lattice QCD

— Bhagwat, Pichowsky, Roberts, Tandy, PRC68, 015203 (03)

## ● Qu-lattice $D_{\text{gluon}}(q)$

Leinweber, Bowman et al  
PRD60, hep-lat/9811027

## ● Find $\Gamma_{\nu}^{\text{eff}}(q, p)$ so DSE produces $S_{\text{latt}}(p)$ from $D_{\text{latt}}(q)$





$$g^2 \gamma_{\mu} D(p - q) Z_{1F}(\mu, \Lambda) \Gamma_{\nu}(q, p) \rightarrow \gamma_{\mu} g^2 D(p - q) \gamma_{\nu} V(p - q)$$

UV limit:  $g^2 D(k^2) V(k^2) \rightarrow \frac{4\pi\alpha_s^{1-\text{loop}}(k^2)}{k^2}$


# Lattice-assisted DSE Results

- Evident vertex enhancement
- Curvature in low  $m_q$  depn
- $M^{\text{IR}}(p^2)$  40% below linear
- Chiral Extrapolation
- $\langle \bar{q}q \rangle_{\mu=1 \text{ GeV}}^{\text{qu-lat}} = -(190 \text{ MeV})^3$
- $\langle \bar{q}q \rangle^{\text{qu-lat}} \approx \langle \bar{q}q \rangle^{\text{expt}} / 2$
- $f_\pi$  30% low




# Qu-lattice $S(p)$ , $D(q)$ mapped to a DSE kernel

$$S(p) = Z(p) [i \not{p} + M(p)]^{-1}$$




# Quenched lattice $\Rightarrow m_q$ Depn of DSE Kernel




Bhagwat, Pichowsky, Roberts, Tandy, PRC68, 015203 (2003)

# Quenched lattice $\Rightarrow m_q$ Depn of DSE Kernel




Bhagwat, Pichowsky, Roberts, Tandy, PRC68, 015203 (2003)

# Quenched lattice $\Rightarrow m_q$ Depn of DSE Kernel




Bhagwat, Pichowsky, Roberts, Tandy, PRC68, 015203 (2003)

# Quenched lattice $\Rightarrow m_q$ Depn of DSE Kernel



Bhagwat, Pichowsky, Roberts, Tandy, PRC68, 015203 (2003)


# Quenched lattice $\Rightarrow m_q$ Depn of DSE Kernel



Bhagwat, Pichowsky, Roberts, Tandy, PRC68, 015203 (2003)


# Quark Confinement—positivity violation

- Confinement/positivity analysis (Osterwalder-Schrader axiom No. 3)
- Fourier transf  $\sigma_S(p_4, \vec{p} = 0)$  to Eucl time  $T$




solid = lattice prop, dashed = MT DSE, dotted = cc pole eg

# DSE and Lattice results for $M_V$ and $M_{ps}$



# Pion electromagnetic form factor

$$\Lambda_\mu = (P' + P)_\mu F_\pi(Q^2) = N_c \int \frac{d^4 q}{(2\pi)^4} \text{Tr} [\bar{\Gamma}^\pi S i\Gamma_\mu S \Gamma^\pi S]$$




# Pion $F(Q^2)$ : Low $Q^2$

(P Maris & PCT, PRC 61, 045202 (2000))

(P. Maris & PCT, PRC 62, 0555204 (2000))


$$r_{\pi}^{\text{DSE}} = 0.68 \text{ fm}$$

$$r_{\pi}^{\text{expt}} = 0.663 \pm .006 \text{ fm}$$




# Kaon $F(Q^2)$ : Low $Q^2$

- Impulse approx + rainbow/ladder  $\Rightarrow$   
conserved em current, correct charge of  $K^+$  and  $K^0$




charge radii	experiment	DSE calc
$r_\pi^2$	$0.44 \pm 0.01 \text{ fm}^2$	$0.45 \text{ fm}^2$
$r_{K^+}^2$	$0.34 \pm 0.05 \text{ fm}^2$	$0.38 \text{ fm}^2$
$r_{K^0}^2$	$-0.054 \pm 0.026 \text{ fm}^2$	$-0.086 \text{ fm}^2$

# Pion electromagnetic form factor




PM and Tandy, PRC62,055204 (2000) [nucl-th/0005015]

# Pion electromagnetic form factor



JLab data from Volmer *et al*, PRL86, 1713 (2001) [nucl-ex/0010009]  
PM and Tandy, PRC62,055204 (2000) [nucl-th/0005015]


# Pion electromagnetic form factor



PM and Tandy, PRC62,055204 (2000) [nucl-th/0005015]


2006a: V. Tadevosyan *et al*, [nucl-ex/0607007], 2006b: T. Horn *et al*, [nucl-ex/0607005]

# 1-loop chiral correction to $r_\pi$ VS $m_\pi$



P. Maris and PCT, in preparation

# 1-loop chiral correction to $r_\pi$ VS $m_\pi$



P. Maris and PCT, in preparation

# LR: Successes, Problems, Resolutions

---

- **Successes:**
  - S-wave mesons, PS and V, light quarks and QQ, no spurious thresholds
  - Exact PS mass formula, Goldstone Thm,  $\Delta M_{HF}$  from DCSB
  - $f_{EW}$ , strong decays, radiative decays, form factors,  $Q^2 < 5\text{GeV}^2$
- **Problems:**
  - Axial vector ( $L > 0$ ) mesons ( $a_1, b_1, \dots$ ) too light
  - Physical diquarks, no physical V or PS  $qQ$  states
  - Excited states are difficult
- **Probable Resolution:**
  - Quark-gluon vertex:  $\Gamma_\mu \Rightarrow \Sigma_q \Rightarrow K_{BSE}$
  - Use analysis of spacelike correlators, 3-pt functions

# From Gluon vertex to BSE Kernel

- A symmetry-preserving procedure [Bender, Roberts, von Smekal, PLB380, (1996), nucl-th/9602012; Munczek 1995] ; Axial vector and vector WTIs, and Goldstone Thm preserved


- $$K_{\text{BSE}}(x', y'; x, y) = -\frac{\delta}{\delta S(x, y)} \Sigma(x', y')$$

- Vertex  $\Gamma_\mu(p, q) = \sum \text{diagrams} \Rightarrow K_{\text{BSE}} = \sum \text{diagrams}$

- If  $\Sigma$  contains:




- $K_{\text{BSE}}$  contains:




- Independent of model parameters. Model does not fight chiral symmetry, use light vector mesons to fix parameters

# $\gamma^* \pi^0 \rightarrow \gamma$ Transition Form Factor




- Abelian axial anomaly +  $\pi$  pole  
in  $\Gamma_{5\mu} \Rightarrow G(0,0)$
- Chiral limit  $G(0,0) = \frac{1}{2}$   
 $\Rightarrow \Gamma_{\pi\gamma\gamma}$  to 2%



# $\gamma^* \pi \gamma^*$ Asymptotic Limit

Lepage and Brodsky, PRD22, 2157 (1980): LC-QCD/OPE  $\Rightarrow$



# Deep Inelastic Lepton Scattering

---

$$W^{\mu\nu} = \frac{1}{2\pi} \text{Disc}_\omega T^{\mu\nu}(\omega) \rightarrow -g^{\mu\nu} \sum_q \frac{e_q^2}{2} f_q(x) + \dots$$

After Bjorken kinematic limit:

$$f_q(x) = \frac{1}{4\pi} \int dz^- e^{iq^+ z^-} \langle \pi(P) | \bar{\psi}_q(\hat{z}) \gamma^+ \psi(0) | \pi(P) \rangle_c$$

$$f_q(x) = \int d^4k \delta(k^+ - x M_\pi) \dots$$

Valence quark number:

$$N_q^V = \int_0^1 dx \{f_q(x) - f_{\bar{q}}(x)\} = \frac{1}{2P^+} \langle \pi(P) | J^+(0) | \pi(P) \rangle = 1$$

# DIS on pion: from DSE-BSE solutions


- Valence quarks, handbag diagram,  $\gamma_\mu$

- Data: J. S. Conway et al, PRD39, 92 (1989)

- Previous: Hecht, Roberts, Schmidt, PRC63, 025213 (2001)


$$\Gamma_\pi(k, P) \approx i\gamma_5 B_0(k^2)/f_\pi^0$$

$S(p)$  fit to data



- Large x behavior:  $(1 - x)^\alpha$ ,  $\alpha = ?$


# DIS on pion: large $x$ behavior?



● Fit:  $a x (1 - x)^{\alpha(x)}$


● BSE ampls: pQCD behavior sets in at a larger scale

# DIS on pion: large $x$ behavior?



- Global fits to (limited) DIS data produce  $\alpha \sim 1.5$
- Parton model (F-J), pQCD (Brodsky, Ezawa), DSEs,  $\Rightarrow \alpha \sim 2+$
- Constituent q models, NJL, duality, etc  $\Rightarrow \alpha \sim 1$

# Quark Distributions in $K$



- Some dependence of  $u(x)$  on environment

# Exact PS Meson Mass Formula for any $m_q$

- Flavor non-singlet and singlet (incl anomaly)

$$\begin{aligned}
 i\partial_\mu(z)\langle j_{5\mu}^\alpha(z)q(x)\bar{q}(y)\rangle &= 2i\mathcal{M}^{\alpha\beta}\langle j_5^\beta(z)q(x)\bar{q}(y)\rangle \\
 &\quad -2i\text{tr}_f(\mathcal{F}^\alpha)\langle Q_t(z)q(x)\bar{q}(y)\rangle \\
 &\quad +\delta(z-x)i\gamma_5\mathcal{F}^\alpha\langle q(x)\bar{q}(y)\rangle + \langle q(x)\bar{q}(y)\rangle i\gamma_5\mathcal{F}^\alpha\delta(y-z)
 \end{aligned}$$

- Matrix elements, amputated  $\Rightarrow$  AV-WTI

$$\begin{aligned}
 P_\mu\Gamma_{5\mu}^\alpha(k;P) &= -2i\mathcal{M}^{\alpha\beta}\Gamma_5^\beta(k;P) -\delta_{\alpha,0}\Gamma_A(k;P) \\
 &\quad +S^{-1}(k_+)i\gamma_5\mathcal{F}^\alpha + i\gamma_5\mathcal{F}^\alpha S^{-1}(k_-)
 \end{aligned}$$

- Residues at PS poles  $\Rightarrow$   $m_p^2 f_p^\alpha = 2\mathcal{M}^{\alpha\beta}\rho_p^\beta + \delta^{\alpha,0}n_p$

$$n_p = 2\text{tr}_f(\mathcal{F}^0)\langle 0|Q_t|p\rangle, \quad \rho_p^\alpha(\mu) = \langle 0|\bar{q}\gamma_5\mathcal{F}^\alpha q|p\rangle, \quad p = \text{any PS}$$

—[Bhagwat, Chang, Liu, Roberts, PCT, PRC 76, 2007; arXiv:0708.1118]

# Chiral limit massless states?

$$\begin{aligned} i\partial_\mu(z) \langle j_{5\mu}^\alpha(z) q(x) \bar{q}(y) \rangle &= 2i \mathcal{M}^{\alpha\beta} \langle j_5^\beta(z) q(x) \bar{q}(y) \rangle \\ &- 2i \text{tr}_f(\mathcal{F}^\alpha) \langle Q_t(z) q(x) \bar{q}(y) \rangle \\ &+ \delta(z-x) i\gamma_5 \mathcal{F}^\alpha \langle q(x) \bar{q}(y) \rangle + \langle q(x) \bar{q}(y) \rangle i\gamma_5 \mathcal{F}^\alpha \delta(y-z) \end{aligned}$$

- Chiral limit, non-singlet flavor:  $\Rightarrow P^2 = 0$  pole forced by DCBS

- No singlet Goldstone boson if, and only if:



$$2i \text{tr}_f(\mathcal{F}^0) \int d^4z \langle Q_t(z) q(0) \bar{q}(0) \rangle \equiv - \int \frac{d^4k}{(2\pi)^4} i\mathcal{F}^0 \{ \gamma_5, S_0(k) \}$$

- That is:  $\boxed{N_f \text{tr}_{\text{cs}} \gamma_5 \langle Q_t q(0) \bar{q}(0) \rangle_\mu \equiv \langle \bar{q} q \rangle_\mu}$

- Can this be verified? Lattice?

# Flavor Non-singlet PS Mass Relation


$$f_H m_H^2 = 2 m_q(\mu) \rho_H(\mu)$$



PM, Roberts, Tandy, PLB420, 267 (1998) [nucl-th/9707003]


# Inaccuracy of GMOR

$qQ$  case:



GMOR: 0.2%( $\pi$ ); 4%(K); 14%(0.4GeV); 30%(D)

# Quark mass functions from DSE solutions




# Constituent Mass Concept for $c$ - and $b$ -quarks

All GeV	D(uc)	D*(uc)	D <sub>s</sub> (sc)	D* <sub>s</sub> (sc)
expt M	1.86	2.01	1.97	2.11
calc M	1.85(FIT)	2.04	1.97	2.17
expt f	0.222	?	0.294	?
calc f	0.154	0.160	0.197	0.180

All GeV	B(ub)	B*(ub)	B <sub>s</sub> (sb)	B* <sub>s</sub> (sb)	B <sub>c</sub> (cb)	B* <sub>c</sub> (cb)
expt M	5.28	5.33	5.37	5.41	6.29	?
calc M	5.27(FIT)	5.32	5.38	5.42	6.36	6.44
expt f	0.176	?	?	?	?	?
calc f	0.105	0.182	0.144	0.20	0.210	0.18

- **Fit**  $\Rightarrow$  constituent masses:  $M_c^{\text{cons}} = 2.0 \text{ GeV}$ ,  $M_b^{\text{cons}} = 5.3 \text{ GeV}$
- Consistent with  $M^{DSE}(p^2 \sim -M^2)$  generated by  $m_c = 1.2 \pm 0.2$ ,  $m_b = 4.2 \pm 0.2$ , [PDG,  $\mu = 2 \text{ GeV}$ ]
- Does heavy quark dressing contribute anything? Too much in this DSE model—no mass shell !

# Compare Quark Masses with PDG



# Quarkonia

All GeV	$M_{\eta_c}$	$f_{\eta_c}$	$M_{J/\psi}$	$f_{J/\psi}$
expt	2.98	0.340	3.09	0.411
calc with $M_c^{\text{cons}}$	3.02	0.239	3.19	0.198
calc with $\Sigma_c^{\text{DSE}}(p^2)$	3.04	0.387	3.24	0.415

All GeV	$M_{\eta_b}$	$f_{\eta_b}$	$M_\Upsilon$	$f_\Upsilon$
expt	9.4 ?	?	9.46	0.708
calc with $M_b^{\text{cons}}$	9.6	0.244	9.65	0.210
calc with $\Sigma_b^{\text{DSE}}(p^2)$	9.59	0.692	9.66	0.682

- QQ and qQ decay constants too low by 30-50% in constituent mass approximation
- Quarkonia decay constants much better for DSE dressed quarks (within 5% of expt.)
- IR sector (gluon  $k$  below  $\sim 0.8$  GeV) contribute little for bb or cc quarkonia in DSE, BSEs
- QQ states are more point-like than qq or qQ states



# Recovery of a $qQ$ Mass Shell

- Suppress gluon  $k$  below  $\sim 0.8$  GeV in DSE dressing of  $b$  propagator
- Retain IR sector for dressed "light" quark and BSE kernel
- Now a mass shell is produced



All GeV	B(ub)	B*(ub)	B <sub>s</sub> (sb)	B* <sub>s</sub> (sb)	B <sub>c</sub> (cb)	B* <sub>c</sub> (cb)
expt M	5.28	5.33	5.37	5.41	6.29	?
calc M	4.66	—	4.75	—	5.83	—
expt f	0.176	?	?	?	?	?
calc f	0.133	—	0.164	—	0.453	—

- Masses are  $\sim 10$  % low
- It makes sense that  $R_b < R_{qQ} \Rightarrow$  greater limit on low  $k$  in  $\Sigma_b$
- May be partial confirmation of Brodsky and Shrock's suggestion of universal maximum wavelength for quarks/gluons in hadrons [Phys. Lett. B666, (2008)]

# IR Suppression of Kernel



# A Schematic Model: Flavor mixing, $\eta, \eta'$



- [Bhagwat, Chang, Liu, Roberts, PCT, PRC (76), 2007; arXiv:0708.1118]
- Structure:  $K_N = \text{LR vector gluon exch}$ ,  
 $K_A = \mathcal{F}(\gamma_5, \not{P}\gamma_5) \otimes (\gamma_5, \not{P}\gamma_5)\mathcal{F}$ ,  $\mathcal{F} = \text{diag}(1/M_f)$
- (Munczek-Nemirovsky) t-channel  $\delta^4(k)$  for  $K_N$  and  $K_A$
- 2 strength parameters:  $\rho^0 \Rightarrow K_N$ ,  $m_{\eta'} \Rightarrow K_A$ .
- Fix  $m_u, m_d, m_s \dots$  via vector mesons

# $\pi^0 - \eta - \eta'$ mixing: 3 flavors

---

- $m_u - m_d$  causes  $\pi^0$  to be mixed in:

$$135 \text{ MeV} : |\pi^0\rangle \sim 0.72 \bar{u}u - 0.69 \bar{d}d - 0.013 \bar{s}s$$

$$455 \text{ MeV} : |\eta\rangle \sim 0.53 \bar{u}u + 0.57 \bar{d}d - 0.63 \bar{s}s$$

$$922 \text{ MeV} : |\eta'\rangle \sim 0.44 \bar{u}u + 0.45 \bar{d}d + 0.78 \bar{s}s$$

- $m_u = m_d \Rightarrow$

$$455 \text{ MeV} : |\eta\rangle \sim 0.55 (\bar{u}u + \bar{d}d) - 0.63 \bar{s}s, \quad \theta_\eta = -15.4^\circ$$

$$924 \text{ MeV} : |\eta'\rangle \sim 0.45 (\bar{u}u + \bar{d}d) + 0.78 \bar{s}s, \quad \theta_{\eta'} = -15.7^\circ$$

- Chiral limit:  $m_{\eta'}^2 = (0.852 \text{ GeV})^2 \equiv 2\text{tr}_f(\mathcal{F}^0) \langle 0|Q_t|\eta'\rangle / f_{\eta'}^0$
- cf Witten-Veneziano a-v ghost scenario  $\Rightarrow m_{\eta'}^2 = h^2 + m_{\text{GB}}^2$
- It is worth extending to a realistic LR model for  $K_N$  with separable  $K_A$ : one obtains access to decay constants, residues, and details of the mass relations

# The V-A Current Correlator

- $\Pi_{\mu\nu}^V(x) = \langle 0 | T j_\mu(x) j_\nu^\dagger(0) | 0 \rangle$  , isovector currents  $j_\mu = \bar{u}\gamma_\mu d$ ,  $j_\mu^5 = \bar{u}\gamma_5\gamma_\mu d$

$$\Pi_{\mu\nu}^V(P) = (P^2\delta_{\mu\nu} - P_\mu P_\nu) \Pi^V(P^2)$$

$$\Pi_{\mu\nu}^A(P) = (P^2\delta_{\mu\nu} - P_\mu P_\nu) \Pi^A(P^2) + P_\mu P_\nu \Pi^L(P^2)$$

$$\Pi_{\mu\nu}^V(P) = - \int_q^\Lambda \gamma_\mu Z_1(\mu, \Lambda) \Gamma_\nu^V(q, P)$$

- $m_q = 0$  :  $\Pi^V - \Pi^A = 0$  , to all orders in pQCD
- $\Pi^V - \Pi^A$  probes the scale for onset of non-perturbative phenomena in QCD

# The 4-quark Condensates

- Operator product expansion  $\Rightarrow$  leading uv behavior

$$\Pi^{V-A}(P^2) = \frac{32\pi\alpha_s \langle \bar{q}q\bar{q}q \rangle}{9 P^6} \left\{ 1 + \frac{\alpha_s}{4\pi} \left[ \frac{247}{4\pi} + \ln\left(\frac{\mu^2}{P^2}\right) \right] \right\} + \mathcal{O}\left(\frac{1}{P^8}\right)$$

- Often **vacuum saturation** ( $\langle \bar{q}q\bar{q}q \rangle \approx \langle \bar{q}q \rangle^2$ ) is assumed for QCD Sum Rules. **Validity not known.**
- Extract  $\langle \bar{q}q\bar{q}q \rangle$  from  $\lim_{P^2 \rightarrow \infty} P^6 \Pi^{V-A}(P^2)$

Model	$-\langle \bar{q}q \rangle_{\mu=19} (GeV)^3$	$\langle \bar{q}q\bar{q}q \rangle_{\mu=19} (GeV)^6$	$R(\mu = 19)$
Set A	$(0.5682)^3$	$(0.619)^6$	1.67
Set B	$(0.1734)^3$	$(0.1902)^6$	1.74
Set C	$(0.2469)^3$	$(0.2695)^6$	1.69
<b>Set D</b>	<b><math>(0.216)^3</math></b>	<b><math>(0.235)^6</math></b>	<b>1.65</b>

—T. Nguyen, PCT, in preparation, 2008

# DSE Calculation: Weinberg Sum Rules

- I:  $\frac{1}{4\pi^2} \int_0^\infty ds [\rho_v(s) - \rho_a(s)] = [P^2 \Pi^{V-A}(P^2)]_{P^2 \rightarrow 0} = -f_\pi^2$
- II:  $P^2 [P^2 \Pi^{V-A}(P^2)]|_{P^2 \rightarrow \infty} = 0$
- DGMLY:  $\int_0^\infty dP^2 [P^2 \Pi^{V-A}(P^2)] = -\frac{4\pi f_\pi^2}{3\alpha} [m_{\pi^\pm}^2 - m_{\pi^0}^2]$

Model	$f_\pi^2$ ( $GeV^2$ )	$f_\pi$ ( $MeV$ )	$f_\pi^{exp} / f_\pi^{num}$	$\Delta m_\pi$ ( $MeV$ )	$(\Delta m_\pi)_{exp}$
Set A	0.00456291	67.5	1.37	4.86	
Set B	0.00538895	73.4	1.26	5.2	$4.43 \pm 0.03$
Set C	0.00518379	72.0	1.28	4.88	

# Summary

---

- Effective ladder-rainbow model based on QCD -DSEs;  $\langle \bar{q}q \rangle_\mu \Rightarrow 1$  IR parameter
- Ground state  $qQ$  and  $QQ$  mesons (V & PS) up to b-quark region
- Constituent mass concept generates good meson masses, but appears inadequate for electroweak decays
- Dynamical dressing in  $S(p)$  at each stage increases the value of the decay constant [factor of 3 for  $\bar{b}b$ , factor of 2 for  $\bar{c}c$ ] !
- Gluon momenta incompatible with hadron size are largely irrelevant for  $cc, bb$  states, but need to be suppressed to even produce  $B, D$  mesons
- First BSE-DSE treatment of pion and kaon DIS for valence  $u(x), s(x)$
- Used  $\langle J J \rangle$ , V-A, to estimate  $\langle \bar{q}q\bar{q}q \rangle$  as  $\sim 70\%$  greater than vac saturation, and npQCD enters at scale 0.5 fm.



# Collaborators

---

- Craig Roberts, Argonne National Lab
- Pieter Maris, Iowa State University
- Yu-xin Liu, Lei Chang, Peking University
- Nick Souchlas, Trang Nguyen, Kent State University

Thankyou!

# Constituent Quark-like Behavior for $c$ , $b$ -quarks



- Mass shell positions marked for  $\bar{b}b$  and  $\bar{c}c$  quarkonia
- $qQ$  mesons sample  $M_Q(p^2) \sim 4$  times further into timelike region
- The same constituent or pole mass is unlikely to suffice for  $QQ$  and  $qQ$  mesons

# General Pseudoscalar Mass Formula

---

- $N_f = 3$ , charge neutral states:  $p = \pi^0, \eta, \eta'$


$$m_p^2 \begin{bmatrix} f_p^3 \\ f_p^8 \\ f_p^0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ n_p \end{bmatrix} + \left[ 2 \mathcal{M}_{3 \times 3} \right] \begin{bmatrix} \rho_p^3 \\ \rho_p^8 \\ \rho_p^0 \end{bmatrix}$$

- Isospin breaking:  $m_u \neq m_d$  allows anomaly,  $\mathcal{F}^0$ , and  $s\bar{s}$  into  $\pi^0$
- $\eta'$  in  $SU(N_f)$  limit:  $m_{\eta'}^2 f_{\eta'}^0 = n_{\eta'} + 2m \rho_{\eta'}^0$

# Model Bethe-Salpeter Kernel for flavor singlet?

- Vertex integral eqns do not involve  $Q_t(x)$  explicitly:  


$$\Gamma_{5\mu}^\alpha(k; P) = Z_2 \gamma_5 \gamma_\mu \mathcal{F}^\alpha + \int^\Lambda K S_+ \Gamma_{5\mu}^\alpha S_-$$
- DSE models need:  $K_{\text{BSE}} = K_{\text{N}} + K_{\text{A}}$ , both are  $\bar{q}q$  irreducible,  $K_{\text{N}}$  is also n-gluon irreducible




- A scenario that works: Witten-Veneziano massless axial-vector ghost linking pseudoscalar GBs

# c- and b-Quark Mass Function for BSE

c,b quark mass function near the peak of the parabolic region with  $P^2$  near the meson mass shells  
 $m_c(19 \text{ GeV})=0.88 \text{ GeV}$ ,  $m_b(19 \text{ GeV})=3.8 \text{ GeV}$



# DSE Calculation: Estimated 4 quark condensate



Model	$-\langle \bar{q}q \rangle_{\mu=19} (\text{GeV})^3$	$\langle \bar{q}q\bar{q}q \rangle_{\mu=19} (\text{GeV})^6$	$R(\mu = 19)$
Set A	$(0.5682)^3$	$(0.619)^6$	1.67
Set B	$(0.1734)^3$	$(0.1902)^6$	1.74
Set C	$(0.2469)^3$	$(0.2695)^6$	1.69
Set D	$(0.216)^3$	$(0.235)^6$	1.65