

# Gluons, quarks and deconfinement at high density

Jon-Ivar Skullerud  
with Simon Hands, Seyong Kim

NUI Maynooth

QCD–TNT, Trento, 9 September 2009

# Outline

## Background

QC<sub>2</sub>D vs QCD

## Formalism

Tensor structures

Lattice formulation

## Results


Bulk thermodynamics

Gluon propagator results

Quark propagator results

## Summary

# Background



- ▶ A plethora of phases at high  $\mu$ , low  $T$
- ▶ Based on models and perturbation theory
- ▶ Details depend on diquark gaps and strange quark mass
- ▶ **Diquark condensation** a generic feature

## Lattice simulations?

A **non-perturbative**, **first-principles** approach is needed!

## Lattice simulations?

A **non-perturbative**, **first-principles** approach is needed!

But **QCD** at  $\mu \neq 0$  has a **sign problem**:

$$\gamma_5 \mathcal{M}(\mu) \gamma_5 = \mathcal{M}^\dagger(-\mu) \implies \det \mathcal{M} \text{ may be complex}$$

So standard Monte Carlo importance sampling can not be used!

## Lattice simulations?

A **non-perturbative**, **first-principles** approach is needed!

But **QCD** at  $\mu \neq 0$  has a **sign problem**:

$$\gamma_5 \mathcal{M}(\mu) \gamma_5 = \mathcal{M}^\dagger(-\mu) \implies \det \mathcal{M} \text{ may be complex}$$

So standard Monte Carlo importance sampling can not be used!

### Indirect approach

Study QCD-like theories without a sign problem

- ▶ **Generic features** of strongly interacting systems at  $\mu \neq 0$
- ▶ Check on **model calculations**

## QC<sub>2</sub>D vs QCD— Issues of interest

**Gluodynamics** — SU(2) and SU(3) very similar?

- ▶ Deconfinement at high density — effects on gluon propagator?
- ▶ Gap equation with effective or one-gluon interaction used to determine superconducting gap → more realistic input?
- ▶ Static magnetic gluon: unscreened at all orders in perturbation theory!

## QC<sub>2</sub>D vs QCD— Issues of interest

**Gluodynamics** — SU(2) and SU(3) very similar?

- ▶ Deconfinement at high density — effects on gluon propagator?
- ▶ Gap equation with effective or one-gluon interaction used to determine superconducting gap → more realistic input?
- ▶ Static magnetic gluon: unscreened at all orders in perturbation theory!

**Quark propagator**

- ▶ Details of phase diagram depend critically on the effective quark mass in the medium.
- ▶ Location of Fermi surface?
- ▶ Direct determination of diquark gap, size of Cooper pairs?

## Tensor structure in medium

The medium breaks Lorentz (Euclidean) symmetry to  $O(3)$

$\Rightarrow$  1  $\rightarrow$  2 scalar functions in gluon, 2  $\rightarrow$  4 in quark:

$$D_{\mu\nu}(\vec{q}, q_t) = P_{\mu\nu}^T D_M(\vec{q}^2, q_t^2) + P_{\mu\nu}^E D_E(\vec{q}^2, q_t^2) + \xi \frac{q_\mu q_\nu}{q^4}$$

$$S^{-1}(\vec{p}, \tilde{\omega}) = i\vec{p} A(\vec{p}^2, \tilde{\omega}^2) + i\gamma_4 \tilde{\omega} C(\vec{p}^2, \tilde{\omega}^2) + B(\vec{p}^2, \tilde{\omega}^2) \\ + i\gamma_4 \vec{p} D(\vec{p}^2, \tilde{\omega}^2)$$

$$S(\vec{p}, \tilde{\omega}) = i\vec{p} S_a + i\gamma_4 \tilde{\omega} S_c + S_b + i\gamma_4 \vec{p} S_d$$

where  $\tilde{\omega} \equiv p_4 - i\mu$ .

## Gor'kov formalism

Quarks and antiquarks are in the same representation.

Construct Gor'kov spinor

$$\Psi = \begin{pmatrix} \psi \\ \bar{\psi}^T \end{pmatrix} \implies \langle \Psi(x) \bar{\Psi}(y) \rangle \equiv \mathcal{G}(x, y) = \begin{pmatrix} S_N & -S_A \\ \bar{S}_A & \bar{S}_N \end{pmatrix}$$

## Gor'kov formalism

Quarks and antiquarks are in the same representation.

Construct Gor'kov spinor

$$\Psi = \begin{pmatrix} \psi \\ \bar{\psi}^T \end{pmatrix} \implies \langle \Psi(x) \bar{\Psi}(y) \rangle \equiv \mathcal{G}(x, y) = \begin{pmatrix} S_N & -S_A \\ \bar{S}_A & \bar{S}_N \end{pmatrix}$$

$S_A$  contains information about anomalous propagation

Self-energies are diquark gaps  $\Delta$  (superfluid/superconducting)

## Gor'kov formalism

Quarks and antiquarks are in the same representation.

Construct Gor'kov spinor

$$\Psi = \begin{pmatrix} \psi \\ \bar{\psi}^T \end{pmatrix} \implies \langle \Psi(x) \bar{\Psi}(y) \rangle \equiv \mathcal{G}(x, y) = \begin{pmatrix} S_N & -S_A \\ \bar{S}_A & \bar{S}_N \end{pmatrix}$$

$S_A$  contains information about anomalous propagation

Self-energies are diquark gaps  $\Delta$  (superfluid/superconducting)

General tensor structure is the same as for normal components.

## Gor'kov formalism

Quarks and antiquarks are in the same representation.

Construct Gor'kov spinor

$$\Psi = \begin{pmatrix} \psi \\ \bar{\psi}^T \end{pmatrix} \implies \langle \Psi(x) \bar{\Psi}(y) \rangle \equiv \mathcal{G}(x, y) = \begin{pmatrix} S_N & -S_A \\ \bar{S}_A & \bar{S}_N \end{pmatrix}$$

$S_A$  contains information about anomalous propagation

Self-energies are diquark gaps  $\Delta$  (superfluid/superconducting)

General tensor structure is the same as for normal components.

### Symmetries

From isospin and charge conjugation symmetry it follows that

$$\bar{S}_N(x, y) = -S_N(y, x)^T, \quad S_A(x, y) = S_A(y, x)^T$$

# Fermi surface and Cooper pairs

## Fermi surface

In a Fermi liquid the Fermi surface is given by

$$\det S^{-1}(\vec{p}_F, p_4 = 0) = 0 \quad \iff \quad \vec{p}^2 A^2 + \tilde{\omega}^2 C^2 + B^2 = 0$$

Pole in propagator

# Fermi surface and Cooper pairs

## Fermi surface

In a Fermi liquid the Fermi surface is given by

$$\det S^{-1}(\vec{p}_F, p_4 = 0) = 0 \quad \iff \quad \vec{p}^2 A^2 + \tilde{\omega}^2 C^2 + B^2 = 0$$

## Pole in propagator

In gapped phase: zero crossing!

# Fermi surface and Cooper pairs

## Fermi surface

In a Fermi liquid the Fermi surface is given by

$$\det S^{-1}(\vec{p}_F, p_4 = 0) = 0 \quad \iff \quad \vec{p}^2 A^2 + \tilde{\omega}^2 C^2 + B^2 = 0$$

## Pole in propagator

In gapped phase: zero crossing!

## Size of Cooper pair

If we know the anomalous propagator  $S_A(x)$  we can compute the size of the Cooper pairs:

$$\xi^2 = \frac{\int d^3x \vec{x}^2 \left| \frac{1}{2} \text{Tr}(S_A(x)\Lambda^+) \right|^2}{\int d^3x \left| \frac{1}{2} \text{Tr}(S_A(x)\Lambda^+) \right|^2}$$

# Lattice formulation

We use **Wilson fermions**:

- ▶ Correct symmetry breaking pattern, Goldstone spectrum
- ▶  $N_f < 4$  needed to guarantee continuum limit
- ▶ No problems with locality, fourth root trick
- ▶ Chiral symmetry buried at bottom of Fermi sea

# Lattice formulation

We use **Wilson fermions**:

- ▶ Correct symmetry breaking pattern, Goldstone spectrum
- ▶  $N_f < 4$  needed to guarantee continuum limit
- ▶ No problems with locality, fourth root trick
- ▶ Chiral symmetry buried at bottom of Fermi sea

$$S = \bar{\psi}_1 M(\mu) \psi_1 + \bar{\psi}_2 M(\mu) \psi_2 - J \bar{\psi}_1 (C \gamma_5) \tau_2 \bar{\psi}_2^T + \bar{J} \psi_2^T (C \gamma_5) \tau_2 \psi_1$$
$$\gamma_5 M(\mu) \gamma_5 = M^\dagger(-\mu), \quad C \gamma_5 \tau_2 M(\mu) C \gamma_5 \tau_2 = -M^*(\mu)$$

# Lattice formulation

We use **Wilson fermions**:

- ▶ Correct symmetry breaking pattern, Goldstone spectrum
- ▶  $N_f < 4$  needed to guarantee continuum limit
- ▶ No problems with locality, fourth root trick
- ▶ Chiral symmetry buried at bottom of Fermi sea

$$S = \bar{\psi}_1 M(\mu) \psi_1 + \bar{\psi}_2 M(\mu) \psi_2 - J \bar{\psi}_1 (C \gamma_5) \tau_2 \bar{\psi}_2^T + \bar{J} \psi_2^T (C \gamma_5) \tau_2 \psi_1$$
$$\gamma_5 M(\mu) \gamma_5 = M^\dagger(-\mu), \quad C \gamma_5 \tau_2 M(\mu) C \gamma_5 \tau_2 = -M^*(\mu)$$

**Diquark source  $J$**  introduced to

- ▶ lift low-lying eigenmodes in the superfluid phase
- ▶ study diquark condensation without uncontrolled approximations

## Simulation Parameters



We work on two lattices, 'coarse' and 'fine'.

Two 'finer' lattices are used for  $\mu = 0$  simulations only

Name	$\beta$	$\kappa$	Volume	$a$	$am_\pi$	$m_\pi/m_\rho$
coarse	1.7	0.178	$8^3 \times 16$	0.23fm	0.79	0.80
fine	1.9	0.168	$12^3 \times 24$	0.18fm	0.65	0.80
finer, h	2.0	0.162	$12^3 \times 24$		0.64	0.83
finer, l	2.0	0.163	$12^3 \times 24$		0.52	0.76


- ▶ Simulations performed with  $j = J/\kappa = 0.04$  for  $\mu = 0.3 - 1.0$
- ▶ 300–500 trajectories for each  $\mu$ .
- ▶ Simulations with  $j = 0.02, 0.06$  for  $\mu = 0.3, 0.5, 0.7, 0.9$  (coarse lattice)  $\rightarrow$  enable extrapolation to  $j = 0$ .

# Thermodynamics results




- ▶ Close to SB scaling for  $\mu > \mu_d$
- ▶ Deconfining transition at  $a\mu_d \sim 0.65$  on **both** lattices?!
- ▶ BEC (superfluid nuclear matter)  $\rightarrow$  BCS (Quarkyonic superfluid) crossover?

## Gluon propagator results





Some finite volume and lattice spacing effects at  $\mu = 0$


In-medium modifications, incl. violations of Lorentz symmetry, visible in magnetic gluon at  $\mu = 0.7$




## Coarse lattice results




Static magnetic gluon extrapolated to  $j=0$ 

Magnetic gluon ( $q_4 = 2\pi T$ ) extrapolated to  $j=0$ 

Electric gluon extrapolated to  $j=0$ 

## Fine lattice results



## In-medium gluon mass

Crude fit to 'massive' form

$$D_{E,M}(\vec{q}, q_4) = \frac{Z}{\vec{q}^2 + q_4^2 + m_{e,m}^2}$$

**not** a good fit!

Improvement:

Try HDL-inspired form?

# In-medium gluon mass


Crude fit to 'massive' form

$$D_{E,M}(\vec{q}, q_4) = \frac{Z}{\vec{q}^2 + q_4^2 + m_{e,m}^2}$$

not a good fit!

Improvement:


Try HDL-inspired form?



Fit gives  $m_e = 0$  for  $a\mu < 0.7$   
on fine lattice

# Volume dependence


[ coarse lattice,  $\mu = 0.9, j = 0.04$  ]



# Quark propagator results

## Quark propagator in vacuum

- ▶ Large lattice spacing dependence
- ▶ Substantial quark mass dependence for  $Z(p)$
- ▶ Unusual  $p$ -dependence in  $Z(p)$
- ▶ infrared suppression recovered in low-mass and continuum limit?
- ▶  $M(p)$  not yet properly corrected!



## Tensor structure

Extracting form factors with the most general Ansatz for the tensor structure is complicated!

We would like to reduce the number of components to consider


# Tensor structure

Extracting form factors with the most general Ansatz for the tensor structure is complicated!


We would like to reduce the number of components to consider

We find that

- ▶ the **vector**, **scalar** and **temporal** components of the **normal** propagator  $S_a, S_b, S_c$  and
- ▶ the **scalar** and **tensor** components  $A_b, A_d$  of the **anomalous** propagator are nonzero
- ▶ all other components are zero

Quark propagator on coarse lattice,  $\mu = 0.5$ 


## Normal propagator: spatial vector part




# Normal propagator: temporal vector part

All data are for  $aj = 0.04$ !


Fermi momentum may be found by extrapolating zero crossing to  $k_4 = 0$




## Normal propagator: scalar part



## Anomalous propagator: scalar part



## Anomalous propagator: tensor part



## Summary

- ▶ Vacuum  $\rightarrow$  BEC  $\rightarrow$  BCS phase

## Summary

- ▶ Vacuum  $\rightarrow$  BEC  $\rightarrow$  BCS phase
- ▶ Magnetic gluon strongly enhanced in BEC phase
- ▶ Screening of both **magnetic** and **electric** gluon propagator in BCS phase
  - $\rightarrow$  Electric: Debye screening
  - $\rightarrow$  Magnetic: Landau damping
  - $\rightarrow$  **Static magnetic** gluon is also screened!

## Summary

- ▶ Vacuum  $\rightarrow$  BEC  $\rightarrow$  BCS phase
- ▶ Magnetic gluon strongly enhanced in BEC phase
- ▶ Screening of both **magnetic** and **electric** gluon propagator in **BCS phase**
  - $\rightarrow$  Electric: Debye screening
  - $\rightarrow$  Magnetic: Landau damping
  - $\rightarrow$  **Static magnetic** gluon is also screened!
- ▶ Strong modifications of quark propagator in BEC phase
  - $\rightarrow$  Zero crossing in **vector** component
    - **evidence of superfluid gap and Fermi surface!**
  - $\rightarrow$  **Scalar component** 'goes away'!
    - chiral condensate rotates into diquark condensate!

## Summary

- ▶ Vacuum  $\rightarrow$  BEC  $\rightarrow$  BCS phase
- ▶ Magnetic gluon strongly enhanced in BEC phase
- ▶ Screening of both **magnetic** and **electric** gluon propagator in BCS phase
  - $\rightarrow$  Electric: Debye screening
  - $\rightarrow$  Magnetic: Landau damping
  - $\rightarrow$  **Static magnetic** gluon is also screened!
- ▶ Strong modifications of quark propagator in BEC phase
  - $\rightarrow$  Zero crossing in **vector** component
    - **evidence of superfluid gap and Fermi surface!**
  - $\rightarrow$  **Scalar component** 'goes away'!
    - chiral condensate rotates into diquark condensate!
- ▶ Clear signal for **anomalous** propagation
  - $\rightarrow$  **Scalar** anomalous propagator becomes  $\approx$  constant at large  $\mu$
  - $\rightarrow$  Need to understand **tensor** component!

## Outlook

- ▶ Extrapolate all results to zero diquark source
- ▶ Invert Gorkov propagator to obtain form factors and gaps
- ▶ Determine Fermi momentum from zero crossing in propagator
- ▶ Determine size of Cooper pairs (BEC→BCS)

# Outlook

- ▶ Extrapolate all results to zero diquark source
- ▶ Invert Gorkov propagator to obtain form factors and gaps
- ▶ Determine Fermi momentum from zero crossing in propagator
- ▶ Determine size of Cooper pairs (BEC→BCS)
- ▶ Study Gribov copy effects / gauge dependence using stereographic Landau gauge [with Dhagash Mehta]
- ▶ Continuum extrapolation?