


Double Helicity Dependence of Jet Properties

Douglas Fields
University of New Mexico

How can you get a net k_T ?

- Nice conceptual picture from D. Sivers (talk given at UNM/RBRC Workshop on Parton Orbital Angular Momentum).
- Summary:
 - Sivers effect requires orbital motion.
 - Is process dependent.
- I would like to just follow this conceptual picture to see where it may lead.




What is the origin of jet k_T ?


$$\frac{\langle p_T^2 \rangle_{\text{pair}}}{2} = \langle k_T^2 \rangle = \langle k_T^2 \rangle_{\text{intrinsic}} + \langle k_T^2 \rangle_{\text{soft}} + \langle k_T^2 \rangle_{\text{NLO}}$$

Intrinsic (Confinement) $k_T \approx 200 \text{ MeV}/c$

Soft QCD radiation.



An example - J/ψ production.



Extra gluon kick

$$\langle p_T \rangle_{J/\psi} = 1.8 \pm 0.23 \pm 0.16 \text{ GeV}/c$$

Phys. Rev. Lett. 92, 051802, (2004).

Origin of Jet k_T

Net transverse momentum of a di-hadron pair in a factorization ansatz

$$\frac{\langle p_T^2 \rangle_{\text{pair}}}{2} = \langle k_T^2 \rangle = \langle k_T^2 \rangle^I \oplus \langle k_T^2 \rangle^S \oplus \langle k_T^2 \rangle^H$$



- $I = \text{initial}$ – fermi motion of the confined partons, initial-state gluon radiation...
- $S = \text{soft}$ – one or several soft gluons emitted
- $H = \text{hard}$ – final-state radiation (three-jet)...

Another Possibility:

- Spin-correlated transverse momentum – partonic orbital angular momentum
- Can examine the helicity dependence of each term:


$$\Delta \langle k_T^2 \rangle_{\text{initial}} \neq 0?$$

$$\Delta \langle k_T^2 \rangle_{\text{soft}} = 0$$

$$\Delta \langle k_T^2 \rangle_{\text{hard}} \neq 0?$$


Mechanism for $\Delta \langle k_T^2 \rangle_{initial} \neq 0$

Same Helicity



Mechanism for $\Delta \langle k_T^2 \rangle_{initial} \neq 0$

Opposite Helicity



Integrating over Impact Parameter

Drell-Yan pair production in polarized $p+p$ collisions.

(Meng Ta-Chung et al. Phys. Rev. D 1989)

$$p_t^2(b, \theta_P, \theta_T; \pm) = \left| \vec{k}_{PR} \right|^2 + \left| \vec{k}_{TR} \right|^2 \pm \left| \vec{k}_{PR} \right| \left| \vec{k}_{TR} \right| \cos \left(\theta_P + \theta_T \right)$$

Averaging over $D(b)$:


$$\Delta \langle p_t^2(b) \rangle = \langle p_t^2(b, +) \rangle - \langle p_t^2(b, -) \rangle$$

Integrating over impact parameter b :


$$\Delta \langle p_t^2 \rangle \approx 1.9 \langle k_R \rangle^2$$

$$\langle k_R \rangle \equiv \langle \left| \vec{k}_{PR} \right| \rangle = \langle \left| \vec{k}_{TR} \right| \rangle$$

Like Helicity



Un-like Helicity





Measuring jet k_T – di-hadron correlation

Intra-jet pairs angular width : $\sigma_{\text{near}} \rightarrow \langle j_T \rangle$

Inter-jet pairs angular width : $\sigma_{\text{far}} \rightarrow \langle j_T \rangle$

- Zero k_T




Measuring jet k_T – di-hadron correlation

Intra-jet pairs angular width : $\sigma_{\text{near}} \rightarrow \langle j_T \rangle$

Inter-jet pairs angular width : $\sigma_{\text{far}} \rightarrow \langle j_T \rangle \oplus \langle k_T \rangle$

- Non-Zero k_T




Jet Kinematics

$$\sqrt{\langle j_T^2 \rangle} = \sqrt{2 \langle j_{Ty}^2 \rangle} \simeq \sqrt{2} \frac{p_{Tt} \cdot p_{Ta}}{\sqrt{p_{Tt}^2 + p_{Ta}^2}} \sigma_N$$

$$\frac{\langle z_t(k_T, x_h) \rangle \sqrt{\langle k_T^2 \rangle}}{\hat{x}_h(k_T, x_h)} = \frac{1}{x_h} \sqrt{\langle p_{out}^2 \rangle - \langle j_{Ty}^2 \rangle} (1 + x_h^2)$$

$$x_h \equiv \frac{p_{Ta}}{p_{Tt}} \quad \hat{x}_h = \hat{x}_h(\langle k_T^2 \rangle, x_h) \equiv \frac{\langle \hat{p}_{Ta} \rangle}{\langle \hat{p}_{Tt} \rangle}$$

$$\langle z_t \rangle = \left\langle \frac{p_{Tt}}{\hat{p}_{Tt}} \right\rangle$$





For details, see PRD 74,
072002 (2006)


Measuring jet k_T – di-hadron correlation

π^0 - h^\pm azimuthal correlation

Trigger on a particle, e.g. π^0 , with transverse momentum p_{Tt} . Measure azimuthal angular distribution *w.r.t.* the azimuth of associated (charged) particle with transverse momentum $p_{T\alpha}$. The strong **same** and **away side peaks** in p-p collisions indicate di-jet origin from hard scattering partons.




PHENIX Detector



•Philosophy:

- High resolution at the cost of acceptance
- High rate capable DAQ
- Excellent **trigger capability** for rare events



•Central Arms:

$$|\eta| < 0.35, \Delta\phi = 2 \times 90^\circ$$

Charged particle ID and tracking;
photon ID. EM Calorimetry.


Muon Arm:

$$1.2 < |\eta| < 2.4$$

Muon ID and tracking.

•Global Detectors

Collision trigger
Collision vertex
characterization
Relative luminosity
Local Polarimetry



π^0 Identification

$$\pi^0 \rightarrow \gamma + \gamma$$



The pions are selected within

$$M_{\pi^0} \pm 2.0\sigma_{\pi^0}$$

Photon trigger:



EMCal cluster energy > 1.4 GeV

The mass, M_{π^0} , and the width, σ_{π^0} , are determined by fitting the di-photon mass spectrum with *gaus+pol3*.



Correlation Function

- **Blue** – real (raw $\Delta\phi$)
- **Red** – background (mixed events)




Fit Function

$$\frac{dN_{real}}{d\Delta\phi} = \frac{1}{N} \frac{dN_{mix}}{d\Delta\phi} \cdot \left(C_o + C_1 \cdot Gaus(0) + C_2 \cdot \frac{dN_{away}}{d\Delta\phi} \Big|_{\pi/2}^{3\pi/2} \right)$$

$$\frac{dN_{away}}{d\Delta\phi} \Big|_{\pi/2}^{3\pi/2} = \frac{-p_{Ta} \cos \Delta\phi}{\sqrt{2\pi} \langle p_{out}^2 \rangle \text{Erf} \left(\sqrt{2} p_{Ta} / \sqrt{\langle p_{out}^2 \rangle} \right)} \exp \left(-\frac{p_{Ta}^2 \sin^2 \Delta\phi}{2 \langle p_{out}^2 \rangle} \right)$$

- Blue – real (raw $\Delta\phi$)
- Red – Fit from above equation





Need $\langle z_t \rangle$ and x_h


$$\frac{\langle z_t(k_T, x_h) \rangle \sqrt{\langle k_T^2 \rangle}}{\hat{x}_h(k_T, x_h)} = \frac{1}{x_h} \sqrt{\langle p_{\text{out}}^2 \rangle - \langle j_{Ty}^2 \rangle (1 + x_h^2)}$$

\hat{x}_h and $\langle z_t \rangle$ are determined through a combined analysis of the π^0 inclusive cross section and using jet fragmentation function measured in e^+e^- collisions. *Ref.* PRD 74, 072002 (2006)

$\langle k_T^2 \rangle$ asymmetry results are not sensitive to this \hat{x}_h and $\langle z_t \rangle$ values – only needed to set the scale.




k_T results for unpolarized case



Preliminary Results

- Take the difference Like – Unlike helicities
- Normalized by beam polarizations
- $\langle j_T^2 \rangle$ asymmetry $(-3 \pm 8(\text{stat}) \pm 5(\text{syst}) \text{ MeV out of } \sim 630 \text{ MeV unpolarized})$
- $\langle k_T^2 \rangle$ asymmetry also small $(-37 \pm 88(\text{stat}) \pm 14(\text{syst}) \text{ MeV out of } \sim 2.8 \text{ GeV unpolarized})$.



Interpretation

- The first term can be related in the “Meng conjecture” to partonic transverse momentum:

$$\Delta \langle k_T^2 \rangle^I = \sum_{i,j} c^{ij} W^{ij} \left\{ \langle \vec{k}_T^i \cdot \vec{k}_T^j \rangle^{++} - \langle \vec{k}_T^i \cdot \vec{k}_T^j \rangle^{+-} \right\}$$

where c^{ij} give the initial state weights and W^{ij} give the impact parameter weighting.

- Since PYTHIA studies show that the final state studied here is dominated by gg scattering (~50%),
- We can interpret this as a constraint on the gluon orbital angular momentum.
- Consistent with previous (Sivers) measurements, and complimentary, since one naïvely needs no initial or final state interactions (although Sivers measures orbital in *transversely* polarized proton).

Relating to Meng...

$$\Delta\langle p_T^2 \rangle \approx 1.9\langle k_\phi \rangle^2$$

yields:

$$\langle k_\phi \rangle \approx \Delta\sqrt{\langle k_T^2 \rangle} = -37 \pm 88^{\text{stat}} \pm 14^{\text{syst}} \text{ MeV}/c,$$

Summary

- We have an analysis tool – di-hadron azimuthal correlations - that allows us to measure k_T .
- We are studying this in helicity-sorted collisions to see if there is a spin-dependent, coherent component of k_T .
- In our kinematic range, we find $\Delta j_T^{RMS} = -3 \pm 8(\text{stat}) \pm 5(\text{syst})$ MeV, and $\Delta k_T^{RMS} = -37 \pm 88 \pm 14(\text{syst})$ MeV.
- While this is consistent with other measurements related to gluon OAM, the connection to parton OAM needs theoretical guidance!