

Model Calculations of T-odd TMDs

TMD2010 - Trento

Aurore Courtoy

INFN- Sezione di Pavia

with S. Scopetta (Perugia) and V. Vento (Valencia)

23/06/2010

Outlines

Motivations

Formalism: The MIT Bag and a Constituent Quark Model

Results


- The Sivers Function

- The Boer-Mulders Function

Bag Model vs. CQM

Conclusions

Prototype Process: Semi-Inclusive Deep Inelastic Scattering




$$\text{SIDIS: } I(l) + N(P) \rightarrow I(l') + h(P_h) + X$$

$Q^2, P \cdot q, P \cdot P_h, P_h \cdot q \rightarrow \infty$ and x, z finite


Factorization

$$W^{\mu\nu} \propto \sum_q e_q^2 \int d^4p d^4k \Phi(k, P, S) \gamma^\mu \Delta(p, P_h) \gamma^\nu$$



- $\Phi(k, P, S) \Rightarrow$ Parton Distribution Functions
- $\Delta(p, P_h) \Rightarrow$ Fragmentation Functions
- Nonperturbative Objects

Some Asymmetries in SIDIS



- ϕ_h = angle between **leptonic** and **hadronic** planes
- ϕ_S = angle between **leptonic** plane and **transverse spin** of the target
- **Trento Convention** [PRD70, 117504]

Azimuthal Asymmetries for unpolarized target in SIDIS

$$\text{e.g., } A(\phi_h) \Rightarrow \langle \cos \phi_h \rangle, \langle \cos 2\phi_h \rangle$$

Single-Spin Asymmetries for transverse target polarization in SIDIS

$$\text{e.g., } A(\phi_h, \phi_S) \Rightarrow \langle \sin(\phi_h - \phi_S) \rangle, \langle \sin(\phi_h + \phi_S) \rangle$$

Transverse Momentum Dependent PDF

Non-perturbative effects of the **intrinsic transverse momentum \vec{k}_\perp of the quarks** inside the nucleon may induce significant hadron azimuthal asymmetries.

[Cahn; Mulders & Tangermans, ...]

- Relaxing Time-reversal Invariance \Rightarrow *naive* **T-odd** functions,


e.g. **Sivers & Boer-Mulders functions**

[Sivers, PRD41];[Boer & Mulders PRD57.]

- Existence of **Final State Interactions** at leading-order

[Brodsky, Hwang & Schmidt, PLB 530];[Belitsky, Ji & Yuan NPB 656.]

- The gauge link:



0th order, No gauge link \longrightarrow T-odd fct = 0

Existence of leading-twist FSI \longrightarrow T-odd fct \neq 0

Models

The Sivers function $f_{1T}^{\perp Q}(x, k_T)$

⇒ Distribution of **unpolarized** quarks inside a **transversely polarized** proton

The Boer-Mulders functions $h_1^{\perp Q}(x, k_T)$

⇒ Distribution of **transversely polarized** quarks inside a **unpolarized** proton



- **non-perturbative** quantities → not calculable in QCD
- we use **models** for the proton → not an exact calculation
- **goal** → insights into microscopic mechanisms
- **HERE:** formalisms for

▶ **MIT bag model**

e.g. [Jaffe, PRD11]

▶ **Constituent Quark Models (CQM)**

e.g. [de Rújula, Georgi & Glashow, PRD12]

Formalisms for the T -odd functionsI. Constituent Quark Model \longrightarrow 3-body

Recipe



- go to a helicity basis [Sivers],
- expand the free quark fields
- properly insert complete sets of free states

- identify the intrinsic proton w.f. $\Psi_r S_z$
- we are left with the interaction term

$$V(\vec{k}_1, \vec{k}_3, \vec{q})$$

$$= \frac{1}{q^2} \bar{u}_{m_1}(\vec{k} - \vec{q}) \Gamma u_{m_2}(\vec{k}) \bar{u}_{m_3}(\vec{k}_3) \gamma^+ u_{m_4}(\vec{k}_3 - \vec{q})$$

\longrightarrow to be reduced NR
(in a CQM fashion)



[A.C., F. Fratini, S. Scopetta and V. Vento, PRD78]

Formalisms for the T -odd functions

II. MIT Bag Model \longrightarrow 1-body

Recipe



- go to a helicity basis [Sivers],
- expand into the **bag quark w.f.**
- properly insert **complete sets of free states**

- we are left with the interaction term

$$V(\vec{k}_1, \vec{k}_3, \vec{q})$$

$$= \frac{1}{q^2} \varphi_{m_1}^\dagger(\vec{k} - \vec{q}) \gamma^0 \Gamma \varphi_{m_2}(\vec{k}) \varphi_{m_3}^\dagger(\vec{k}_3) \gamma^0 \gamma^+ \varphi_{m_4}(\vec{k}_3 - \vec{q})$$

[F. Yuan, PLB575]



The interaction

I. Constituent Quark Model

NR reduction of the interaction - up to $O\left(\frac{k^2}{m^2}\right)$ -

Use of **free spinors** \longrightarrow

$$u_m(\vec{k}) \propto \begin{pmatrix} \chi_m \\ \frac{\vec{\sigma} \cdot \vec{k}}{k^0 + m} \chi_m \end{pmatrix}$$

$f_{1T}^{\perp Q}, h_1^{\perp Q} \neq 0$ comes from **Interference of the lower and upper components** in the four-spinors of the **free quark states**

The interaction is to be calculated between **proton states** in a CQM \Rightarrow e.g., **Harmonic Oscillator** $|N\rangle = a^{|2} S_{1/2}\rangle_S$

\Rightarrow **SU(6)** symmetry for the proton

II. MIT Bag Model

Bag wave function \longrightarrow

$$\varphi_m(\vec{k}) \propto \begin{pmatrix} t_0(|\vec{k}|) \chi_m \\ \vec{\sigma} \cdot \hat{k} t_1(|\vec{k}|) \chi_m \end{pmatrix}$$

$f_{1T}^{\perp Q}, h_1^{\perp Q} \neq 0$ comes from the **Interference of the lower and upper components** in the **bag w.f.**

The interaction is to be calculated between **proton states**, we choose \Rightarrow **SU(6)** symmetry for the proton

Properties of the Sivers function

Experiment

- ▶ Evidence for **non-zero** Sivers function at HERMES [2003]
- ▶ Sivers Asymmetry **statistically compatible with zero** within present statistical error at COMPASS
- ▶ **Future**: CLAS@12GeV?

Extraction from data

- ▶ W. Vogelsang and F. Yuan, PRD 72, 054028 (2005)
- ▶ M. Anselmino et al., Eur.Phys.J.A39:89-100,2009
- ▶ J.C. Collins et al., PRD 73, 014021 (2006)

Theory: Properties of the Sivers function

- ▶ From first principles: **Burkardt Sum Rule** (PRD 69 (2004) 091501)

$$\sum_{Q=u,d} \langle k_x^Q \rangle = \sum_{Q=u,d} - \int_0^1 dx \int d\vec{k}_T \frac{k_x^2}{M} f_{1T}^{\perp Q}(x, k_T) = 0$$

- ▶ Hypothetical relation with the E GPD
 - distribution for u is **negative**,
 - distribution for d is **positive**.
- ▶ other **Model Calculations**


Model Calculations with SU(6) proton WF and perturbative OGE

NR Constituent Quark Model

[A.C., Fratini, Scopetta and Vento, PRD 78]

- ▶ 3-body calculation
- ▶ No proportionality u and d distribution
- ▶ **Small Violation** of the Burkardt SR

$$\frac{\langle k_x^u \rangle + \langle k_x^d \rangle}{\langle k_x^u \rangle - \langle k_x^d \rangle} \simeq 0.02$$




MIT Bag Model



[A.C., Scopetta and Vento, PRD 79]

- ▶ 1-body calculation
- ▶ No proportionality u and d distribution
- ▶ **Small Violation** of the Burkardt SR

$$\frac{\langle k_x^u \rangle + \langle k_x^d \rangle}{\langle k_x^u \rangle - \langle k_x^d \rangle} \simeq 0.05$$




Helicity-flip contributions



1st moment



$$f_{1T}^{\perp(1)q}(x) = \int d^2\vec{k}_T \frac{k_T^2}{2M^2} f_{1T}^{\perp q}(x, k_T)$$

- u upper; d lower panels
- **red curves**: results in a CQM (H.O.)
- **blue curves**: results in the MIT bag model
- **full curves**: full results
- **dashed curves**: results interacting quark flipping helicity



Both **Models** at the hadronic scale $\mu_0^2 \sim 0.1 \text{ GeV}^2$

Comparison with extracted Sivers functions: e.g. [Collins et al., PRD73]



1st moment

$$f_{1T}^{\perp(1)q}(x) = \int d^2\vec{k}_T \frac{k_T^2}{2M^2} f_{1T}^{\perp q}(x, k_T) .$$

- **full**: results at the **hadronic scale** $\mu_o^2 \simeq 0.1 \text{ GeV}^2$
- **shaded area**: 1- σ region of the best fit of the Sivers function extracted from HERMES data, at $Q^2 = 2.5 \text{ GeV}^2$

Model at $\sim 0.1 \text{ GeV}^2$ vs. **Exp.** at 2.5 GeV^2

- **Evolution**
 - ▶ **Blue**: results after *NLO-standard* evolution to $Q^2 = 2.5 \text{ GeV}^2$
 - ▶ **Correct evolution missing**

The results in the CQM and the MIT bag Model are \sim in agreement with the extraction of the Sivers function by Collins et al. and by Anselmino et al.

Properties of the Boer-Mulders function

● Experiment

- ▶ Drell-Yan at FNAL [2007]
- ▶ SIDIS at COMPASS and HERMES [2009]
- ▶ **Future:** PAX@FAIR ?
- ▶ Difficulty of the analysis

● Extraction from data

- Drell-Yan** [Zhang, Lu, Ma and Schmidt, PRD 77]
SIDIS predictions [Barone, Prokudin and Ma, PRD78]
both [Barone, Prokudin and Melis, 0912.5194]

● Theory: Properties of the Boer-Mulders function


- ▶ Hypothetical relation with the chiral-odd \bar{E} GPD [Burkardt and Hannafious, PLB658]
 - distribution for u is negative,
 - distribution for d is negative.
- ▶ From first principles: **Lattice** [QCDSF and UKQCD Colls., PRL 98]
 - moments of chiral-odd GPDs
- ▶ other **Model Calculations**

Model Calculations with SU(6) proton WF and perturbative OGE

NR Constituent Quark Model

[A.C., Scopetta and Vento, PRD 80]


- ▶ 3-body calculation
- ▶ no-proportionality u and d distributions





MIT Bag Model

[A.C., Scopetta and Vento, PRD 80]

- ▶ 1-body calculation
- ▶ no-proportionality u and d distributions




Helicity-flip contributions



1st moment


$$h_1^{\perp(1)q}(x) = \int d^2\vec{k}_T \frac{k_T^2}{2M^2} h_1^{\perp q}(x, k_T).$$

- u upper; d lower panels
- **red curves:** results in a CQM (H.O.)
- **blue curves:** results in the MIT bag model
- **full curves:** full results
- **dashed curves:** results non-flipping helicity



Both Models at the hadronic scale $\mu_0^2 \sim 0.1 \text{ GeV}^2$

Comparison with extracted BM functions: e.g. [Barone et al., 0912.5194]





1st moment

$$h_1^{\perp(1)q}(x) = \int d^2\vec{k}_T \frac{k_T^2}{2M^2} h_1^{\perp q}(x, k_T) .$$

- **full**: results for d distributions
- **dashed**: results for u distributions
- **MIT Bag** at the hadronic scale $\mu_o^2 \simeq 0.1 \text{ GeV}^2$
- **NR CQM** at the hadronic scale $\mu_o^2 \simeq 0.1 \text{ GeV}^2$
- **parametrization** from HERMES and COMPASS data, at $Q^2 \sim 2.5 \text{ GeV}^2$
- **Evolution**
 - ▶ I should evolve the results to $Q^2 = 2.5 \text{ GeV}^2$ (like we did for the Sivers function)
 - ▶ **Correct** evolution **missing** so I haven't done it yet . . .

Comparison of the T -odd functions



⇒ Boer-Mulders bigger than Sivers function for both flavors

⇒ same trend in both models for both flavors.

Conclusions for the CQM

T -odd functions in CQM:

- Analysis of the Siverts & Boer-Mulders functions in a 3-Body model
[A.C., Fratini, Scopetta, Vento, PRD78]
- Formalism valid for any CQM
⇒ **Ingredients**: wave functions and a reduction of the interaction
- Non-Relativistic

Siverts function in the H.O.

[A.C., Fratini, Scopetta, Vento, PRD78]

- ▶ Correct relative **sign** for u and d distributions
- ▶ **Burkardt sum rule** recovered
- ▶ **Reasonable agreement** with extraction from data

Boer-Mulders function in the H.O.

[A.C., Scopetta, Vento, PRD80]

- ▶ Correct relative **sign** for u and d distributions
- ▶ **Reasonable agreement** with expectations from other evaluations

Conclusions for the MIT Bag Model

T -odd functions in the Bag:

- Analysis of the Sivers & Boer-Mulders functions in a 1-Body model [F. Yuan, PLB575]
- Formalism in the bag
 ⇒ **Ingredients**: bag wave functions and SU(6) proton state
- Relativistic

Sivers function in the **bag**

[A.C., Scopetta, Vento, PRD79]

- ▶ Correct relative **sign** for u and d distributions
- ▶ **Burkardt sum rule** recovered
- ▶ **Reasonable agreement** with extraction from data

Boer-Mulders function in the **bag**

[A.C., Scopetta, Vento, PRD80]

- ▶ Correct relative **sign** for u and d distributions
- ▶ **Reasonable agreement** with expectations from other evaluations

Conclusions


Conclusions

- **Agreement between models**
 - ▶ More *physical* picture for **helicity-flip at the quark level**
 ⇒ either the *active* and *non-active* quark can flip
 - ▶ First Principles arguments ⇒ **Burkardt Sum Rule** fulfilled [**Sivers**]
- Much **better agreement** of the MIT Bag calculation with the **current knowledge**
- **By-Product:** more confidence on the **Non-Relativistic** expansion.
- **... to the experiments**
 1. Need for Correct **evolution** of TMDs
 talks of Cherednikov, Manashov, ...
 2. Relation between **GPDs in Impact Parameter Space** and T -odd functions?
 see talk of Matthias Burkardt, Barbara Pasquini, Cédric Lorcé, ...
 3. **Improvement of the extractions** ; need for more experimental data
 talk of Stefano Melis, Marco Radici, ...
 4. **Improvement of the models** after experimental feedback.
LOOP: experimentalists could use updated model calculations!

**I don't think we did go blind,
I think we are blind,
Blind but seeing,
Blind people who can see, but do not see.**

Jose Saramago

Calculation Details: MIT bag



$$f_{1T}^{\perp Q}(x, k_{\perp}) \propto 2\Re \left\{ \int \frac{d^2 q_{\perp}}{(2\pi)^5} \frac{i}{q^2} \sum_{\{m\}, \beta} C_{\{m\}}^{\mathcal{Q}, \beta} \varphi_{m_1}^{\dagger}(\vec{k} - \vec{q}_{\perp}) \gamma^0 \gamma^+ \varphi_{m_2}(\vec{k}) \int \frac{d^3 k_3}{(2\pi)^3} \varphi_{m_3}^{\dagger}(\vec{k}_3) \gamma^0 \gamma^+ \varphi_{m_4}(\vec{k}_3 - \vec{q}_{\perp}) \right\}$$

$$\int \frac{d^3 k_3}{(2\pi)^3} \varphi_{m_3}^{\dagger}(\vec{k}_3) \gamma^0 \gamma^+ \varphi_{m_4}(\vec{k}_3 - \vec{q}_{\perp}) = F(\vec{q}_{\perp}) \delta_{m_3 m_4} + H(\vec{q}_{\perp}) \delta_{m_3, -m_4}$$


With the MIT Bag WF,

$$\varphi_m(\vec{k}) \propto \begin{pmatrix} t_0(|\vec{k}|) \chi_m \\ \vec{\sigma} \cdot \vec{k} t_1(|\vec{k}|) \chi_m \end{pmatrix}, \quad t_i(k) = \int_0^1 u^2 du j_i(ukR_0) j_i(u\omega)$$

$H(\vec{q}_{\perp})$ does not vanish in a basis for the gluon's momentum constrained by the DIS framework,
 \Rightarrow z-axis is the virtual photon's direction
 \Rightarrow operator structure: γ^+

Sivers function in a CQM: Calculation details


In a helicity basis, to the first non-vanishing order
by expanding the **free quark fields**
and by properly inserting **complete sets of free states**



$$\begin{aligned}
 f_{1T}^{\perp Q}(\mathbf{x}, \mathbf{k}_T) &= \Im \left\{ \frac{M}{2k_x} \int \frac{d\xi^- d^2\vec{\xi}_T}{(2\pi)^3} e^{-i(x\xi^- P^+ - \vec{\xi}_T \cdot \vec{k}_T)} \langle PrS_z = 1 | \right. \\
 &\int d\vec{k}_3 \sum_{m_3} b_{m_3 i}^{\mathcal{Q}^\dagger}(\vec{k}_3) e^{ik_3^+ \xi^- - i\vec{k}_3 T \cdot \vec{\xi}_T} \bar{u}_{m_3}(\vec{k}_3) \\
 &\sum_{l_n, l_1} \int d\vec{k}_n \int d\vec{k}_1 |\vec{k}_1 l_1\rangle |\vec{k}_n l_n\rangle \langle \vec{k}_n l_n | \langle \vec{k}_1 l_1 | \\
 &(ig) \int_{\xi^-}^{\infty} A_a^+(0, \eta^-, \vec{\xi}_T) d\eta^- T_{ij}^a \\
 &\sum_{l'_n, l'_1} \int d\vec{k}'_n \int d\vec{k}'_1 |\vec{k}'_1 l'_1\rangle |\vec{k}'_n l'_n\rangle \langle \vec{k}'_n l'_n | \langle \vec{k}'_1 l'_1 | \gamma^+ \\
 &\left. \sum_{m'_3} \int d\vec{k}'_3 b_{m'_3 j}^{\mathcal{Q}}(\vec{k}'_3) u_{m'_3}(\vec{k}'_3) |PS_z = -1\rangle + \text{h.c.} \right\} .
 \end{aligned}$$

[A.C., F. Fratini, S. Scopetta and V. Vento, Phys.Rev.D78:034002,2008.]

Sivers function in a CQM: Calculation



Identifying the **intrinsic proton wave function**:

$$\begin{aligned}
 \mathbf{f}_{1T}^{\perp Q}(\mathbf{x}, \mathbf{k}_T) &= \Im \left\{ i g^2 \frac{M}{2k_x} \int d\vec{k}_1 d\vec{k}_3 \frac{d^4 q}{(2\pi)^3} \delta(q^+) (2\pi) \delta(q_0) \delta(k_3^+ + q^+ - xP^+) \delta(\vec{k}_3 T + \vec{q} T - \vec{k} T) \right. \\
 &\quad \sum_{\mathcal{F}_1, \{m_j\} \{c_j\}} \Psi_{r S_z=1}^\dagger \left(\vec{k}_3 \{m_3, i, Q\}; \vec{k}_1 \{m_1, c_1, \mathcal{F}_1\}; \vec{P} - \vec{k}_3 - \vec{k}_1, l_n \right) T_{ij}^a T_{c_1 c_1'}^a \frac{1}{q^2} V(\vec{k}_1, \vec{k}_3, \vec{q}) \\
 &\quad \left. \Psi_{r S_z=-1} \left(\vec{k}_3 + \vec{q}, \{m_3', j, Q\}; \vec{k}_1 - \vec{q}, \{m_1', c_1', \mathcal{F}_1\}; \vec{P} - \vec{k}_3 - \vec{k}_1, l_n \right) \right\}
 \end{aligned}$$

with the **interaction** given by:

$$V(\vec{k}_1, \vec{k}_3, \vec{q}) = \bar{u}_{m_3}(\vec{k}_3) \gamma^+ u_{m_3'}(\vec{k}_3 + \vec{q}) \bar{u}_{m_1}(\vec{k}_1) \gamma^+ u_{m_1'}(\vec{k}_1 - \vec{q})$$

Next step: reduction of the interaction (in a CQM fashion)

[de Rújula, Georgi, Glashow PRD 12, 147, (1975)]

Calculation details: CQM

NR reduction of the interaction - up to $O\left(\frac{k^2}{m^2}\right)$ -

Use of **free spinors** \longrightarrow

$$u_m(\vec{k}) \propto \begin{pmatrix} \chi \\ \frac{\vec{\sigma} \cdot \vec{k}}{k^0 + m} \chi \end{pmatrix}$$


$$\begin{aligned} V(\vec{k}_1, \vec{k}_3, \vec{q}) &= \left\{ -i \frac{(\vec{q} \times \vec{\sigma}_1)_z}{4m^2} \left(1 + \frac{k_3^z}{m} + \frac{\vec{q} \cdot \vec{k}_3}{4m^2} \right) + i \frac{(\vec{q} \times \vec{\sigma}_3)_z}{2m} \left(1 + \frac{k_1^z}{m} - \frac{\vec{q} \cdot \vec{k}_1}{4m^2} \right) \right. \\ &+ \frac{\vec{\sigma}_3 \cdot (\vec{k}_3 \times \vec{q})(\vec{q} \times \vec{\sigma}_1)_z}{8m^3} + \frac{(\vec{q} \times \vec{\sigma}_3)_z \vec{\sigma}_1 \cdot (\vec{k}_1 \times \vec{q})}{8m^3} \\ &\left. + i \frac{\vec{\sigma}_3 \cdot (\vec{k}_3 \times \vec{q})}{4m^2} - i \frac{\vec{\sigma}_1 \cdot (\vec{k}_1 \times \vec{q})}{4m^2} + O\left(\frac{k_1^2}{m^2}, \frac{k_3^2}{m^2}\right) \right\} \end{aligned}$$

- **helicity-flip** interaction $\rightarrow f_{1T}^{\perp Q}(x, k_T) \neq 0$
- **extreme NR limit** \rightarrow no "small components" of the four-spinors \rightarrow **no helicity-flip**


$f_{1T}^{\perp Q}(x, k_T) \neq 0$ comes from the **Interference of the "small" and "large" components** in the four-spinors of the free quark states

The interaction is to be calculated between **proton states** $\Psi_r S_z = \pm 1$ in a CQM \Rightarrow e.g., **Isgur-Karl**

Sivers function in Isgur-Karl: Higher waves decomposition



total S — S' — M — S-S' — M-S — M-S' —



- **Nucleon state:** (we use the 3 first waves)

$$|N\rangle = a|^2S_{1/2}\rangle_S + b|^2S'_{1/2}\rangle_S + c|^2S_{1/2}\rangle_M$$

Notation: $|^{2S+1}X_J\rangle_t$; $t = A, M, S =$ symmetry type
From spectroscopy:

$$a = 0.933, b = -0.275, c = -0.233$$

- **"Higher waves"**
 - ▶ Importance of small components in the proton wave function
 - ▶ relevance of further analysis with other (relativistic) models.

Spin dependence of the matrix element: Siverts function

In a helicity basis, the matrix element to be evaluated is of the type



$$\begin{aligned}
 & 3 \langle \psi(\vec{k}) \varphi_c \frac{1}{\sqrt{2}} (\phi_{MA} \chi_{MA}^\uparrow + \phi_{MS} \chi_{MS}^\uparrow) | \frac{1 \pm \tau_3(3)}{2} \hat{O}_{spin}(\vec{k}) | \psi(\vec{k}) \varphi_c \frac{1}{\sqrt{2}} (\phi_{MA} \chi_{MA}^\downarrow + \phi_{MS} \chi_{MS}^\downarrow) \rangle \\
 = & 3 \left(-\frac{2}{3} \right) \frac{1}{2} \left\{ \phi_{MA}^* \frac{1 \pm \tau_3(3)}{2} \phi_{MA} \langle \psi(\vec{k}) \chi_{MA}^\uparrow | \hat{O}_{spin}(\vec{k}) | \psi(\vec{k}) \chi_{MA}^\downarrow \rangle \right. \\
 & \left. + \phi_{MS}^* \frac{1 \pm \tau_3(3)}{2} \phi_{MS} \langle \psi(\vec{k}) \chi_{MS}^\uparrow | \hat{O}_{spin}(\vec{k}) | \psi(\vec{k}) \chi_{MS}^\downarrow \rangle + 0 + 0 \right\}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{u} \Rightarrow & 3 \left(-\frac{2}{3} \right) \frac{1}{2} \left\{ 1 \langle \psi(\vec{k}) \chi_{MA}^\uparrow | \hat{O}_{spin}(\vec{k}) | \psi(\vec{k}) \chi_{MA}^\downarrow \rangle + \frac{1}{3} \langle \psi(\vec{k}) \chi_{MS}^\uparrow | \hat{O}_{spin}(\vec{k}) | \psi(\vec{k}) \chi_{MS}^\downarrow \rangle \right\} \\
 = & - \left(f(\vec{k}) + \frac{1}{3} g(\vec{k}) \right)
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{d} \Rightarrow & 3 \left(-\frac{2}{3} \right) \frac{1}{2} \left\{ 0 \langle \psi(\vec{k}) \chi_{MA}^\uparrow | \hat{O}_{spin}(\vec{k}) | \psi(\vec{k}) \chi_{MA}^\downarrow \rangle + \frac{2}{3} \langle \psi(\vec{k}) \chi_{MS}^\uparrow | \hat{O}_{spin}(\vec{k}) | \psi(\vec{k}) \chi_{MS}^\downarrow \rangle \right\} \\
 = & - \left(\frac{2}{3} g(\vec{k}) \right)
 \end{aligned}$$

No proportionality between the u and d -distributions due to the spin and momentum dependence of the operator!


Comparison of the T -odd functions



⇒ Boer-Mulders bigger than Sivers function for both flavors

⇒ same trend in both models for both flavors.

Results in a CQM



1st moment

$$f_{1T}^{\perp(1)q}(x) = \int d^2\vec{k}_T \frac{k_T^2}{2M^2} f_{1T}^{\perp q}(x, k_T) .$$


- **full**: results at the **hadronic scale** $\mu_o^2 \simeq 0.1 \text{ GeV}^2$
- **Shaded Area**: Siverson function extracted from HERMES and COMPASS data
[M. Anselmino et al., Eur. Phys. J. A39:89-100 (2009)]

Model at $\sim 0.1 \text{ GeV}^2$ vs. **Exp.** at 2.5 GeV^2

- **Evolution**
 - ▶ **Blue**: results after *NLO-standard* evolution to $Q^2 = 2.5 \text{ GeV}^2$
 - ▶ **Correct** evolution **missing**

The results in the CQM are \sim in agreement with this extraction of the Siverson function

Revised Results in the MIT Bag Model




- **Shaded Area:** $1-\sigma$ region of the best fit of the Sievers function extracted from HERMES data, at $Q^2 = 2.5$ GeV 2
[Collins et al., PRD 73 (2006) 014021]
- **Dashed Curve:** 1st result in the MIT Bag model *after NLO-standard evolution*
[Yuan, PLB 575 (2003)]
- **Plain blue Curve:** revised result in the MIT Bag model *after NLO-standard evolution*
[A. C., Scopetta & Vento, PRD79]

The results in the MIT Bag Model are in agreement with this extraction of the Sievers function

... up to correct Evolution of the Sievers function

Revised Results in the MIT Bag Model



- **Shaded Area:** **Sivers function extracted from HERMES and COMPASS data**
[M. Anselmino et al., *Eur. Phys. J. A39:89-100 (2009)*]
- **Dashed Curve:** **1st result in the MIT Bag model after NLO-standard evolution**
[Yuan, *PLB 575 (2003)*]
- **Plain blue Curve:** **revised result in the MIT Bag model after NLO-standard evolution**
[A. C., Scopetta & Vento, arXiv:0811.1191 [hep-ph]]

The results in the MIT Bag Model are now in a better agreement with this extraction of the Sivers function

... up to correct Evolution of the Sivers function

Definitions

The Sivers function

Distribution of **unpolarized quarks** inside a **transversely polarized proton**

$$\begin{aligned}
 f_{1T}^{\perp Q}(x, \mathbf{k}_T) &= f_{q/p\uparrow}^Q(x, \vec{\mathbf{k}}_T, \mathbf{S}) - f_{q/p\downarrow}^Q(x, \vec{\mathbf{k}}_T, \mathbf{S}) \\
 &= -\frac{M}{2k_x} \int \frac{d\xi^- d^2\vec{\xi}_T}{(2\pi)^3} e^{-i(x\xi^- P^+ - \vec{\xi}_T \cdot \vec{\mathbf{k}}_T)} \\
 &\quad \frac{1}{2} \sum_{S_y = -1, 1} S_y \langle P, S_y | \bar{\psi}_Q(0, \xi^-, \vec{\xi}_T) \mathcal{L}_{\xi_T}^{\dagger}(\infty, \xi^-) \gamma^+ \mathcal{L}_0(\infty, 0) \psi_Q(0, 0, 0) | P, S_y \rangle
 \end{aligned}$$

The Boer-Mulders function

Distribution of **transversely polarized quarks** inside a **unpolarized proton**

$$\begin{aligned}
 h_1^{\perp Q}(x, \mathbf{k}_T) &= f_{q\uparrow/p}^Q(x, \vec{\mathbf{k}}_T, \mathbf{S}) - f_{q\downarrow/p}^Q(x, \vec{\mathbf{k}}_T, \mathbf{S}) \\
 &= -\frac{M}{2k_x} \int \frac{d\xi^- d^2\vec{\xi}_T}{(2\pi)^3} e^{-i(x\xi^- P^+ - \vec{\xi}_T \cdot \vec{\mathbf{k}}_T)} \\
 &\quad \frac{1}{2} \sum_{S_z = -1, 1} \langle P, S_z | \bar{\psi}_Q(0, \xi^-, \vec{\xi}_T) \mathcal{L}_{\xi_T}^{\dagger}(\infty, \xi^-) \gamma^+ \gamma^2 \gamma_5 \mathcal{L}_0(\infty, 0) \psi_Q(0, 0, 0) | P, S_z \rangle
 \end{aligned}$$